Mount Auburn Hospital
Doctor's Office Building - Suite 511
300 Mount Auburn Street
Cambridge, MA 02238
617-876-1600 \* Fax 617-547-3792

BETH ISRAEL DEACONESS HOSPITAL - MILTON DOCTOR'S OFFICE BUILDING - SUITE 226 100 HIGHLAND STREET MILTON, MA 02186 617-696-3200 ° FAX 617-696-0766

## GUY M. ROCHMAN, M.D.

Assistant Professor of Surgery Harvard Medical School



## **WOUND CARE INSTRUCTIONS**

- The purpose is to keep your suture line clean and free of crusts. This will promote better wound healing.
- 2. The supplies you will need are hydrogen peroxide, gauze sponges, and Bacitracin ointment unless instructed otherwise. These supplies are available in the first aid section of the pharmacy.
- 3. Beginning the day following your procedure, gently remove the dressing.
- 4. Gently cleanse the stitches with a gauze sponge moistened with peroxide. The purpose is to prevent crusts and scabs from forming on suture line.
- 5. Apply a small amount of Bacitracin ointment to the incision.
- 6. This procedure is to be done 2-3 times a day until your next office appointment.
- 7. Keep the wound dry. Avoid washing, showering the area until given permission to do by Dr. Rochman.
- 8. If you notice increased pain, redness, swelling or bleeding at the wound site, please contact the office immediately.
- 9. You need not apply another dressing unless instructed to do so.

10.	Other instructions: